




Figure S2. standard sigmoidal growth curve that assumes 3% N in final biomass used for nutrient dosing above (Table S1).

Experiment 1 Chlorophyll Fluorescence Statistical Analysis

Treatment: 100% NS, 75% NS, 50% NS, 25% NS

Nutrient solution (NS)

Table S1. (A,B,C,D,E): Experiment 1 chlorophyll fluorescence statistical analysis.

Two Way Anova Table						
(A)						
Trait	Term	df	SS	MS	F	P
Alpha (Fv/Fo)	Treatment	3	0.003090	0.001030	5.23	0.002730
Alpha (Fv/Fo)	DAP	7	0.002560	0.000366	1.86	0.091200
Alpha (Fv/Fo)	Treatment:DAP	21	0.002910	0.000139	0.704	0.813000
Alpha (Fv/Fo)	Residual	64	0.012600	0.000197	NA	NA
(B)						
95% TUKEY HSD, Alpha (Fv/Fo)						
100%NS	75%NS	50%NS	25%NS			
0.85 ± 0.01 a	0.84 ± 0.01 ab	0.84 ± 0.02 a	0.83 ± 0.01 b			
(C)						
Trait	Term	df	SS	MS	F	P
Y (Fm)	Treatment	3	0.002770	0.000924	2.62	0.0581
Y (Fm)	DAP	7	0.027400	0.003920	11.1	4.10 × 10 ⁻⁹
Y (Fm)	Treatment:DAP	21	0.004480	0.000213	0.606	0.899
Y (Fm)	Residual	64	0.022500	0.000352	NA	NA
(D)						
Trait	Term	df	SS	MS	F	P
ETR	Treatment	3	747.000000	249.000000	26.0	4.06 × 10 ⁻¹¹
ETR	DAP	7	3476.000000	497.000000	51.8	5.74 × 10 ⁻²⁴
ETR	Treatment:DAP	21	779.000000	37.100000	3.87	1.49 × 10 ⁻⁵
ETR	Residual	64	613.000000	9.580000	NA	NA
(E)						
95% TUKEY HSD, ETR						
DAP	100%NS	75%NS	50%NS	25%NS		
24	19.87 ± 1.26 ab	17.97 ± 1.80 b	22.27 ± 1.36 a	17.87 ± 1.02 b		
26	18.50 ± 1.31 a	13.87 ± 1.27 b	17.43 ± 1.10 a	11.87 ± 0.51 b		
28	26.30 ± 2.36 a	22.37 ± 0.85 a	26.33 ± 2.12 a	20.77 ± 3.07 a		
31	12.77 ± 2.11 a	12.10 ± 2.04 a	13.27 ± 1.62 a	13.07 ± 1.90 a		
33	41.40 ± 5.14 a	22.53 ± 1.79 b	25.33 ± 2.48 b	23.63 ± 8.80 b		
35	16.73 ± 1.88 a	10.70 ± 3.82 ab	6.87 ± 1.34 b	11.90 ± 1.40 ab		
38	33.37 ± 3.79 a	17.47 ± 6.65 b	20.67 ± 3.43 b	21.67 ± 4.18 ab		
39	14.30 ± 1.78 a	9.63 ± 1.31 a	11.03 ± 5.37 a	10.27 ± 1.50 a		

Different letters (a and b) in each row represent significant differences ($p < 0.05$). Chlorophyll fluorescence parameters of basil (*Ocimum basilicum* cv Genovese) measured under different % Sonneveld's solution ($n = 3$): effective PSII quantum yield (Y(Fm)), a proxy for PSII efficiency (Alpha (Fv/Fo)), and electron transport rate (ETR, $\mu\text{mol e}^- \text{m}^{-2} \text{s}^{-1}$).

Experiment 2 Chlorophyll Fluorescence Statistical Analysis

Treatment: 80% NS, 60% NS, 40% NS, 20% NS

Nutrient solution (NS)

Table S2. (A,B,C,D,E,F): Experiment 2 chlorophyll fluorescence statistical analysis.

TWO WAY ANOVA TABLE						
(A)						
Trait	Term	df	SS	MS	F	P
Alpha (Fv/Fo)	Treatment	3	0.0217	0.00723	15.0	1.11×10^{-6}
Alpha (Fv/Fo)	DAP	4	0.00166	0.000414	0.857	0.498
Alpha (Fv/Fo)	Treatment:DAP	12	0.00969	0.000807	1.67	0.111
Alpha (Fv/Fo)	Residual	40	0.0193	0.000483	NA	NA
(B)						
95% TUKEY HSD, Alpha (Fv/Fo)						
80%NS	60%NS	40%NS		20%NS		
0.84 ± 0.01 a	0.79 ± 0.04 b	0.83 ± 0.01 a		0.83 ± 0.01 a		
(C)						
Trait	Term	df	SS	MS	F	P
Y (Fm)	Treatment	3	0.0465	0.0155	22.2	1.29×10^{-8}
Y (Fm)	DAP	4	0.0772	0.0193	27.6	4.91×10^{-11}
Y (Fm)	Treatment:DAP	12	0.0196	0.00163	2.33	0.0226
Y (Fm)	Residual	40	0.028	0.000699	NA	NA
(D)						
95% TUKEY HSD, Y (Fm)						
DAP	80%NS	60%NS	40%NS		20%NS	
31	0.65 ± 0.03 a	0.56 ± 0.02 b	0.64 ± 0.03 ab		0.61 ± 0.05 ab	
33	0.70 ± 0.02 a	0.63 ± 0.02 b	0.69 ± 0.01 a		0.69 ± 0.01 a	
35	0.72 ± 0.01 a	0.58 ± 0.07 b	0.67 ± 0.02 ab		0.71 ± 0.00 a	
38	0.74 ± 0.01 a	0.70 ± 0.04 a	0.74 ± 0.01 a		0.74 ± 0.01 a	
40	0.69 ± 0.03 a	0.67 ± 0.02 a	0.68 ± 0.01 a		0.69 ± 0.01 a	
(E)						
Trait	Term	df	SS	MS	F	P
ETR	Treatment	3	1649	550	3.24	0.032
ETR	DAP	4	14,559	3640	21.5	1.63×10^{-9}
ETR	Treatment:DAP	12	1472	123	0.723	0.721
ETR	Residual	40	6787	170	NA	NA
(F)						
95% TUKEY HSD, ETR						
80%NS	60%NS	40%NS		20%NS		
25.49 ± 17.13 b	39.10 ± 26.48 a	28.03 ± 17.00 ab		33.49 ± 18.61 ab		

Different letters (a and b) in each row represent significant differences ($p < 0.05$). Chlorophyll fluorescence parameters of basil (*Ocimum basilicum* cv Genovese) measured under different % Sonneveld's solution (n = 3): effective PSII quantum yield (Y(Fm)), a proxy for PSII efficiency (Alpha (Fv/Fo)), and electron transport rate (ETR, $\mu\text{mol e}^- \text{m}^{-2} \text{s}^{-1}$).

Experiment 3: Chlorophyll Fluorescence Statistical Analysis

Treatment: 100% NS, 50% NS + 50% WW, 50% NS + 50% WW + 0.46 mM Gypsum

Nutrient solution (NS)

Table S3. (A,B,C,D,E,F): Experiment 3 chlorophyll fluorescence statistical analysis.

TWO WAY ANOVA TABLE						
(A)						
95% TUKEY HSD, Alpha (Fv/Fo)						
Trait	Term	df	SS	MS	F	P
Alpha (Fv/Fo)	Treatment	2	0.00923	0.00462	9.41	0.00042
Alpha (Fv/Fo)	DAP	6	0.00493	0.000822	1.68	0.151
Alpha (Fv/Fo)	Treatment:DAP	12	0.00641	0.000534	1.09	0.393
Alpha (Fv/Fo)	Residual	42	0.0206	0.00049	NA	NA

Table S3. Cont.

TWO WAY ANOVA TABLE						
(B)						
95% TUKEY HSD, Alpha (Fv/Fo)						
100% NS	50% NS + 50% WW		50% NS + 50% WW + Gypsum			
0.86 ± 0.02 a	0.83 ± 0.03 b		0.84 ± 0.02 b			
(C)						
Trait	Term	df	SS	MS	F	P
Y (Fm)	Treatment	2	0.00137	0.000687	0.545	0.584
Y (Fm)	DAP	6	0.0292	0.00486	3.86	0.00375
Y (Fm)	Treatment:DAP	12	0.0392	0.00327	2.59	0.0113
Y (Fm)	Residual	42	0.0529	0.00126	NA	NA
(D)						
95% TUKEY HSD, Y (Fm)						
DAP	100% NS	50% NS + 50% WW		50% NS + 50% WW + Gypsum		
21	0.73 ± 0.01 a	0.70 ± 0.01 c		0.72 ± 0.01 b		
25	0.70 ± 0.02 a	0.67 ± 0.03 a		0.66 ± 0.02 a		
27	0.65 ± 0.04 a	0.66 ± 0.04 a		0.69 ± 0.00 a		
29	0.72 ± 0.06 a	0.73 ± 0.01 a		0.73 ± 0.02 a		
32	0.66 ± 0.08 a	0.71 ± 0.01 a		0.65 ± 0.05 a		
34	0.68 ± 0.07 a	0.73 ± 0.02 a		0.70 ± 0.04 a		
36	0.73 ± 0.02 a	0.61 ± 0.05 b		0.69 ± 0.02 ab		
(E)						
Trait	Term	df	SS	MS	F	P
ETR	Treatment	2	35.2	17.6	3.42	0.0421
ETR	DAP	6	1309	218	42.3	2.96 × 10 ⁻¹⁶
ETR	Treatment:DAP	12	283	23.6	4.57	1.09 × 10 ⁻⁴
ETR	Residual	42	216	5.15	NA	NA
(F)						
95% TUKEY HSD, ETR						
DAP	100% NS	50% NS + 50% WW		50% NS + 50% WW + Gypsum		
21	6.93 ± 0.42 ab	8.00 ± 1.04 a		6.00 ± 0.26 b		
25	14.20 ± 5.54 a	17.23 ± 1.65 a		19.87 ± 4.95 a		
27	17.17 ± 1.36 a	15.03 ± 2.42 a		18.25 ± 1.95 a		
29	8.17 ± 0.59 ab	6.07 ± 1.36 b		8.70 ± 0.17 a		
32	16.50 ± 2.26 a	17.13 ± 3.26 a		13.60 ± 0.35 a		
34	17.60 ± 2.36 a	13.67 ± 1.85 ab		12.60 ± 1.15 b		
36	19.57 ± 0.12 b	14.70 ± 1.22 c		25.43 ± 2.76 a		

Different letters (a, b and c) in each row represent significant differences ($p < 0.05$). Chlorophyll fluorescence parameters of basil (*Ocimum basilicum* cv Genovese) measured under different % Sonneveld's solution, with/without catholyte substitution (n = 3): effective PSII quantum yield (Y(Fm)), a proxy for PSII efficiency (Alpha (Fv/Fo), and electron transport rate (ETR, $\mu\text{mol e}^- \text{m}^{-2} \text{s}^{-1}$).